

Reduced point mass or multipole base functions.

C.C.Tscherning, M.Veicherts,¹ M.Herceg²,

¹*Niels Bohr Institute, University of Copenhagen, Denmark.*

²*DTU Space, Denmark.*

Juliane Maries Vej 30, DK 2100 Copenhagen Oe, Denmark.

Abstract: Point-mass functions or multipole base-functions are harmonic functions, which may be used to represent the (anomalous) gravity potential (T) globally or locally. The functions may be expressed by closed expressions or as sums of Legendre series. In both cases at least the two first terms must be removed since they are not present in T . For local applications the effect of a global gravity model is generally removed (and later restored). Then more terms need to be removed or substituted by terms similar to error-degree variances. We have done some calculations to illustrate the effect of reducing the point mass or multipole functions, i.e. showing how the first zero-crossing as a function of spherical distance comes closer to zero when more terms are removed.

1. Introduction.

Linear combinations of point mass functions or mass multipoles have been used for the representation of the global (W) or regional anomalous gravity potential, T , see e.g. Balmino (1974), Hauck and Lelgemann (1985), Vermeer (1982, 1989, 1990,1992,1993), Marchenko et al., (2001), Ballani et al. (1993), Wu (1984).

The anomalous potential T , is equal to the difference between W and a global gravity field model like EGM96 (Lemoine et al., 1998), ie. it is a harmonic function.

For a point mass base function we have for an approximation to T :

$$\tilde{T} = \sum_{i=1}^l GM_i / l_i \quad (1)$$

where G is the gravitational constant, M_i is the mass, I the number of point masses and l_i is the distance from the mass located at the point Q_i to the point of evaluation, P , see Fig. 1. The distance from the origin to P and Q_i is denoted r_P , r_{Q_i} , respectively and the first will always be larger than the other. The angle (spherical distance) between the vectors to P and Q_i is denoted ψ .




Fig. 1.

We now drop the subscript on Q . The distance l , from P to Q is then $l = \sqrt{r_P^2 + r_Q^2 - 2r_P r_Q \cos(\psi)}$. For the inverse distance we have

$$\frac{1}{l} = \frac{1}{r_P} \sum_{i=0}^{\infty} \left(\frac{r_Q}{r_P}\right)^i P_i(\cos\psi) \quad (2)$$

where P_i are the Legendre polynomials. Multipole-functions are derive from the inverse distance function by integration or differentiation (compare Tscherning and Rapp (1974)), and will be denoted f .

$$\tilde{T}(P) = \sum_{i=1}^M \alpha_i f_i(P) \quad (3)$$

The estimate \tilde{T} is determined so that $\sum_{i=1}^M (\{L_i \tilde{T}\} - obs_i)^2 = \min$, where M is the number of observations and L_i are the linear functionals associated with the observations. In the computations described below we only consider height anomalies (geoid heights), gravity disturbances and second-order radial derivatives.

We intend to show that in order to make the functions suitable for regional gravity field modeling low-degree terms may be removed or substituted by appropriate weights.

2. Higher order derivatives,

For derivatives with respect to r_p we have series expansions similar to eq. (2), where the terms for the first derivative are multiplied with $-(i+1)/r_p$ and for the second derivative with $(i+1)*(i+2)/r_p^2$. Closed expressions for the derivatives are easily found.

$$\frac{\partial \frac{1}{l}}{\partial r_p} = \frac{(r_Q t - r_p)}{l^3} \quad (4)$$

$$\frac{\partial^2 \frac{1}{l}}{\partial r_p^2} = (3(r_p - r_Q t)^2/l^2 - 1)/l^3 \quad (5)$$

Point masses are not the only harmonic functions which may be used as base functions when approximating T , see e.g. Tscherning (1972), Hauck and Lelgemann (1985) but we will only deal with point mass and excentric multipoles base functions, since they fully represents the message of this paper.

3. Reduced point masses.

From eq. (2) we see that if we use the representation eq. (1), it will contain terms of degree zero and one, which are not present in T . So they have to be removed, simply by subtracting from the closed expressions the first two terms in eq. (2) or its derivative. (This is not done for the examples of closed expressions in section 5). But what if we subtract from the data (and later add back) the contribution from a global model like EGM96, complete to degree N ?

$$\tilde{T}_N = \frac{GM}{r_p} \sum_{n=0}^N \left(\frac{a}{r_p}\right)^n \sum_{m=-n}^n C_{nm} P_{nm}(\sin\varphi) \begin{cases} \cos(m\lambda), m \geq 0 \\ \sin(|m|\lambda), m < 0 \end{cases} \quad (6)$$

Here φ is the geocentric latitude, λ the longitude, P_{nm} the normalized Legendre functions and a a scale factor close to the semi-major axis. C_{nm} are the normalized Stokes coefficients with error-estimates σ_{nm}^2 . Degree-variances and error-degree-variances are sums of the C_{nm} squared, σ_{nm}^2 respectively, for a fixed degree, n , multiplied with $(GM/a)^2$.

As pointed out by Arabelos (1980) we can not simply put to zero the first N terms. Here a solution was found, i.e. that the first terms were not put to zero, but put equal to the so-called error-degree variances, $\sigma_{e,i}^2$ contingently scaled by a factor α so as to reflect if the model was better ($\alpha \leq 1$) or worse ($\alpha \geq 1$) in an area.

To get a little more insight into this, let us interpret the point mass potential as a reproducing kernel in a Hilbert space, where the functions are harmonic down to a Bjerhammar-sphere with radius R_B inside the Earth and $r_Q < R_B$. Using a Kelvin transformation we obtain a point D outside the sphere

$$\frac{R_B^2}{r_P r_D} = \frac{r_Q}{r_P} \quad (7)$$

So that $L = \sqrt{1 - 2 \frac{R_B^2}{r_P r_D} \cos(\psi) + \left(\frac{R_B^2}{r_P r_D}\right)^2}$. Then $K(P,D) = \frac{R_B^2}{r_P r_D} / L$ is the so-called

Krarup kernel, Krarup (1969). When interpreted as a covariance function, it has unitless degree-variances equal to 1, i.e. it is not well suited to represent the anomalous potential, since the degree-variances of T tends to zero like $n^{-3} * q$, $q < 1$, see Tscherning and Rapp (1974). So instead of point mass base functions one should consider using potentials of other types of masses, like a bar, see Hauck and Lelgemann (1985, Fig. 2) or excentric multipoles (Marchenko et al., 2001).

There are other inherent problems using mass-type base functions. Which depth should be used for the masses. Should they form a grid? (Vermeer (1990)). This has been studied extensively by e.g. Barthelmes and Kautzleben (1983) and Barthelmes (1985). We will not discuss this here, but show tables of reduced point mass functions.

4. Excentric multipoles – covariance functions.

The covariance functions generally used in least-squares collocation may be interpreted as excentric multipoles using the Kelvin transformation. For two points outside the Earth, P and D, we have typically (Tscherning and Rapp, (1974)) for the covariance of the anomalous potential,

$$cov(r_P, r_D, \psi) = \sum_{i=2}^{\infty} \frac{A}{(i-1)(i-2)(i+4)} \left(\frac{R_B^2}{r_D r_P}\right)^{i+1} P_i(\cos\psi) \quad (8)$$

which is the linear combination of 3 excentric multipoles:

$$F_k(r_p, r_Q, \psi) = \sum_{i=2}^{\infty} \frac{A_k}{(i-k)} \left(\frac{r_Q}{r_p}\right)^{i+1} P_i(\cos\psi) \quad (9)$$

Each of these may be represented by a closed expression. The first terms corresponding to the maximal degree of the global model used, should be substituted by the (scaled potential) error-degree-variances.

$$cov_N(r_p, r_D, \psi) = \sum_{i=2}^N a\sigma_i^e \left(\frac{R_B^2}{r_D r_p}\right)^{i+1} P_i(\cos\psi) \quad (10)$$

5. Reduced point mass and excentric multipole base functions - examples.

The interesting functions are those which represent the geoid (height anomaly) and the first and second derivative with respect to r_p . All quantities are *anomalous* quantities with respect to EGM96 complete to degree $N=24$. We consider geoid and gravity disturbance at height zero and the second order radial derivative at 250 km altitude. For the latter we will put the mass-point at depth 242.5 km, corresponding to data (r_p) at 250 km. The Bjerhammar sphere will be put at 1.46 km depth.

In the first example (Table 1) we use as observation **one** (anomalous) radial gravity gradient observation equal to 0.15 Eötvös, at height 250 km ($M=1$ in eq. (3)). The latitude and the longitude are set to zero for the mass-point, but the latitude for the point P increases in steps from zero. Table 2 shows the same function, now with the quantities evaluated at zero altitude.

In Table 3 we have corresponding values obtained using the covariance function eq. (8), (10) and again with the same gravity gradient observation as used in Table 1 and 2.

We have here calculated values with spherical distance 1.0 degree and so that the value for the second order radial derivative is 0.15 Eötvös.

Table 1. Closed and Reduced Point mass functions at altitude 250 km. Note the location of the first zero-crossing. For the reduced function all terms up to degree 24 have been set equal to zero.

ψ	Closed expression	Reduced to deg. 24.	Closed	Reduced	Closed	Reduced
	Geoid (m)		Gravity disturbance (mgal)		2. order radial deriv. (Eötvös)	
0.0	0.85	0.29	-2.41	-1.79	0.150	0.150
1.0	0.80	0.24	-2.03	-1.37	0.106	0.100
2.0	0.70	0.14	-1.34	-0.62	0.043	0.028
3.0	0.59	0.05	-0.81	-0.09	0.012	-0.005
4.0	0.49	-0.02	-0.49	0.16	0.002	-0.014
5.0	0.42	-0.05	-0.31	0.24	-0.001	0.013
6.0	0.37	-0.05	-0.21	0.23	-0.002	-0.010
7.0	0.32	-0.05	-0.14	0.17	-0.002	-0.007
8.0	0.29	-0.03	-0.11	0.10	-0.001	-0.003
9.0	0.26	-0.01	-0.10	0.01	-0.001	0.000
10.0	0.23	0.00	-0.06	-0.02	0.000	0.002

Table 2. Closed and Reduced Point mass functions at zero altitude. Note the location of the first zero-crossing and the values for zero spherical distance. The location is for the geoid at 3^0 . This corresponds (approximately) to the location of the first zero-point for the first Legendre polynomial in the series eq. (8). (Legendre polynomials have as many zero points as the degree in the interval from -90^0 to 90^0 distributed approximately equidistantly, see Heiskanen and Moritz, 1967, Fig. 1-8).

ψ	Closed expression	Reduced to deg. 24.	Closed	Reduced	Closed	Reduced
	Geoid (m)		Gravity disturbance (mgal)		2. order radial deriv. (Eötvös)	
0.0	3.56	3.13	-49.2	-55.3	13.9	16.1
1.0	1.93	1.24	-7.87	-7.26	-0.11	-0.18
2.0	1.09	0.31	-1.48	0.04	-0.14	-0.21
3.0	0.75	0.02	-0.51	0.98	-0.053	-0.10

Table 3. Values computed using a degree-variance model (eq.(8)) with depth to the Bjerhammar-sphere equal to 1.56 km. Error.degree-variances from EGM96 used to degree 24, scaled with $\alpha= 1.03$. Computed using Least-Squares-Collocation with only one observation (completely equivalent to using eq. (3)).

ψ	Second order radial derivative. Altitude= 250 km. Eötvöes units.	Geoid. Zero altitude. Units m.	Gravity anomaly. Zero altitude. Units mgal.
0.0	0.15	2.11	13.11
1.0	0.13	1.80	10.36
2.0	0.08	1.23	5.33
3.0	0.03	0.46	1.40
4.0	-0.03	-0.04	-0.87
5.0	-0.03	-0.32	-1.86
6.0	-0.03	-0.43	-2.01
7.0	-0.02	-0.41	-1.66
8.0	-0.02	-0.30	-1.08

Again we see the location of the first zero-crossing, which makes the function well suited for representing data where the contribution from EGM96 to degree 24 has been subtracted.

A FORTRAN program redpmass.f for doing these or similar calculations is available as <http://cct.gfy.ku.dk/redpmass.f> , 2010.05.20.

4. Conclusion.

Closed or reduced point mass or multipole functions may be used to represent the anomalous potential. When used regionally referring to a global gravity field model, the first terms must be removed or substituted by error-degree-variances.

For point-mass or multipole functions the terms up to the lowest degree of the reference potential (the global model) have here been put equal to zero. However, it might be possible to find (unitless) terms representing the power in the

frequencies which the global model have not removed, corresponding to error-degree variances.

If covariance functions (corresponding to multipole base functions) are used, error degree-variances (scaled) may be used. This assures that the model in an appropriate manner weights the regional frequencies with respect to the global model used.

References:

- Arabelos, D.: Untersuchungen zur gravimetrischen Geoidbestimmung, dargestellt am Testgebiet Griechenland. Wiss. Arb. d. Fachrichtung Vermessungswesen d. Universitaet Hannover, Hannover 1980.
- Ballani, L., J.Engel and E.Grafarend: Global base functions for the mass density in the interior of a massive body (Earth), Manuscripta Geodaetica, Vol. 18, no. 2, pp. 99-114, 1993.
- Balmino, G.: La Representation du Potentielle par Masses Ponctuelles. Bulletin Geodesique, No. 3, 1974.
- Barthelmes,F.: Untersuchungen zur Approximation des aeusseren Gravitationsfeldes der Erde durch Punktmassen mit optimierten Positionen. Veroeff. Zentralinstituts fuer Physik der Erde, Nr. 92,Potsdam, DDR, 1986.
- Barthelmes,F. and H. Kautzleben: A New Method of Modelling the Gravity Field of the Earth by Point Masses. Proc. of the IAG Symposia, Vol.1, pp 442-448, Dep. of Geodetic Science and Surveying, The Ohio State University, 1983.
- Hauck, H. and D.Lelgemann: Regional gravity field approximation with buried masses using least-norm collocation. Manuscripta Geodaetica, Vol. 10, no. 1, pp. 50-58, 1985.
- Heiskanen, W.A. and H. Moritz: Physical Geodesy. W.H. Freeman & Co, San Francisco, 1967.
- Krarp, T.: A Contribution to the Mathematical Foundation of Physical Geodesy. Meddelelse no. 44, Geodætisk Institut, København 1969.
- Lemoine, F.G., S.C. Kenyon, J.K. Factor, R.G. Trimmer, N.K. Pavlis, D.S. Chinn, C.M. Cox, S.M. Klosko, S.B. Luthcke, M.H. Torrence, Y.M. Wang, R.G. Williamson, E.C. Pavlis, R.H. Rapp, and T.R. Olson, The Development of the Joint NASA GSFC and the National Imagery and Mapping Agency (NIMA) Geopotential Model EGM96, NASA/TP-1998-206861, Goddard Space Flight Center, Greenbelt, MD, July, 1998.
- Marchenko, A.N., F.Barthelmes, U.Meyer and P.Schwintzer: Regional geoid determination: An application to airborne gravity data in the Skagerrak. STR01/07, GFZ Potsdam, 2001.

- Tscherning, C.C. and R.H.Rapp: Closed Covariance Expressions for Gravity Anomalies, Geoid Undulations, and Deflections of the Vertical Implied by Anomaly Degree-Variance Models. Reports of the Department of Geodetic Science No. 208, The Ohio State University, Columbus, Ohio, 1974.
- Tscherning, C.C.: Representation of Covariance Functions Related to the Anomalous Potential of the Earth using Reproducing Kernels. The Danish Geodetic Institute Internal Report No. 3, 1972.
- Vermeer, M.: The use of mass point models for describing the Finnish gravity field. Proceedings, 9th Meeting Nordic Geodetic Commission, Sept. 1982, Gävle, Sweden 1982.
- Vermeer, M.: Modelling of gravity gradient tensor observations using buried shells of mass quadrupoles. In: Lecture Notes for Nordiska Forskarkurser 18/1988 "Modern Techniques in Geodesy and Surveying", Ebeltoft, Denmark, Sept. 1988, pp. 461-478. National Survey and Cadastre of Denmark, Publications 4. Series Vol. 1, Copenhagen 1989.
- Vermeer, M.: Work on satellite gravity gradiometry using a buried masses grid representation. Study on Precise Gravity Field Determination Methods and mission requirements (Phase 2), Final Report, Part 510B, pp. 1-18, ESA contract 8153/88/F/FL, 1990
- Vermeer, M.: FGI Studies on Satellite Gravity Gradiometry. 1. Experiments with a 5-degree buried masses representation. Report 89:3, Finnish Geodetic Institute, Helsinki, 1989.
- Vermeer, M.: Geoid determination with mass point frequency domain inversion in the Mediterranean. Mare Nostrum 2, GEOMED report, Madrid, pp. 109-119, 1992.
- Vermeer, M.: Optimal combination of satellite gravity gradiometry and satellite tracking by GPS: Simulations using frequency domain mass points inversion software. 19 pp. In: Study of the gravity field determination using gradiometry and GPS, Phase 1, Final report ESA Contract 9877/92/F/FL, April 1993.
- Wu, X.: Point-mass model of local gravity field. Acta Geod. et Cartog. Vol.13, 1984.