

# Computer Assignment – Part 1

## Linear inversion through Tikhonov regularization

A vertical fault separates two areas of the subsurface. The first area is a homogeneous quarter space (to the left of the fault in Figure 1) with everywhere the same density  $\rho = 2600 \text{ kg/m}^3$ . On the other side of the fault the density varies with depth  $z$  (positive downwards). Points where the gravity is measured are indicated by triangles in Figure 1.




Figure 1: Outline of a vertical fault model.

The right column in the files "gravdata-n.txt" ( $n = 1, 2, 3, \dots$ ) contains the horizontal gravity gradient (in units of  $s^{-2}$ ) measured at 18 points along a linear, horizontal profile, starting at the fault, on which it is perpendicular. The left column contains the corresponding  $x$ -coordinates in units of  $km$  for the 18 observation points. It is assumed that the vertical fault is located at  $x = 0 \text{ km}$ .

If  $\Delta\rho(z)$  is the vertical density variation in the quarter space on the right, minus the constant density in the quarter space on the left, the horizontal gravity gradient, as a function of  $x$ , is theoretically given by

$$d_j = \frac{\partial g}{\partial x}(x_j) = \int_0^{\infty} \frac{2Gz}{x_j^2 + z^2} \Delta\rho(z) dz$$

where  $G$  is the Gravity constant and  $x_j$  is the horizontal coordinate of the  $j$ 'th observation, measured from the fault plane.

### The inverse problem

We wish to estimate  $\Delta\rho(z)$  from observations of the horizontal gravity gradient along the  $x$ -axis. The first step of the analysis is a discretization of the problem.

- Perform such a discretization by representing the subsurface to the right of the fault by 100 horizontal layers of thickness  $1 \text{ km}$  each. The 100 layers are resting on an infinite

quarter space of density  $2600 \text{ kg/m}^3$  everywhere. It can be shown that the contribution to data from a homogeneous “half layer” (one of the layers in the layer series to the right of the fault, after the discretization) is

$$G\Delta\rho \log \left( \frac{z_{base}^2 + x^2}{z_{top}^2 + x^2} \right),$$

where  $z_{top}$  is the depth to the top of the homogeneous half layer,  $z_{base}$  is the depth to the base of the layer,  $\Delta\rho$  is the density contrast (to the material to the left of the half layer), and  $x$  is the horizontal distance to the edge of the half layer.

- Show that the problem is linear (and hence can be written  $\mathbf{d} = \mathbf{G}\mathbf{m}$ ).
- Is the solution to this problem unique? Why/why not?
- Data is measured with an uncertainty of  $\pm 1.0 \cdot 10^{-9} s^{-2}$ . Find a solution to the problem using *Tikhonov Regularization*. The solution should be physically acceptable, give the best possible resolution of the model, and at the same time fit the data “within its uncertainties”.
- Describe the resolution of the inverse operator used.
- Compute the corresponding uncertainty of the solution.